

UNCLASSIFIED

AD _ 401 566 _

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

401,566

(5) 17511

411496

S/033/62/039/005/010/011
E032/E314

STEP

AUTHOR: ⑧ Saakyan, R.A.

TITLE: ⑥ Additional acceleration in the motion of celestial bodies

PERIODICAL: ④ Astronomicheskiy zhurnal, v. 39, 5, 1962,

pp. 931 - 937

TEXT: It is pointed out that when the centre of mass of a system of two bodies executes a curvilinear motion an additional acceleration appears in the equation for the relative motion of the two bodies. This additional acceleration is given by:

$$S = 2[\vec{U}_r \vec{\omega}] + [\vec{\omega} [\vec{a} \vec{\omega}]] + [\vec{a} \frac{d\vec{\omega}}{dt}] \quad (8)$$

where $\vec{\omega}$ is the angular velocity of the centre of gravity, \vec{U}_r/dt is the velocity of relative motion in the centre-of-mass system, \vec{a} is the distance between the two bodies and the square brackets represent vector products. This additional acceleration is said to be a real effect, independent of the choice of the coordinate system. It gives rise to a force

Card 1/2

Additional acceleration

S/033/62/039/005/010/011
E032/E314

which must be added to the gravitational attraction in order to obtain a correct equation for the relative motion of the two bodies. It immediately follows that the fact that the Sun moves around the centre of the galaxy, and the centre of the galaxy itself may also be in a curvilinear motion, must be taken into account in studying the motion of planets relative to the Sun and the motion of satellites relative to planets. In fact, this is a general conclusion for all binary systems. In particular, the relative motion of the Moon-Earth system, which was solved by Laplace and Hill, is subject to this correction, but the latter has not so far been taken into account. There are 2 tables.

ASSOCIATION: Byurakanskaya astrofizicheskaya observatoriya
Akademii nauk ArmSSR (Byurakan Astrophysical
Observatory, AS ArmSSR)

SUBMITTED: July 5, 1961

Card 2/2

ASTROPHYSICAL COPY